arXiv:1609.07162v1 [math.NT] 22 Sep 2016

A NOTE ON PERMUTATION BINOMIALS AND TRINOMIALS OVER FINITE FIELDS

NERANGA FERNANDO

ABSTRACT. Let p be a prime and e be a positive integer. We completely explain the permutation binomials and trinomials arising from the reversed Dickson polynomials of the $(k+1)$-th kind $D_{n,k}(1,x)$ over $\mathbb{F}_{p^e}$ when $n = p^l + 2$, where $l \in \mathbb{N}$.

1. INTRODUCTION

Let p be a prime and e be a positive integer. Let $\mathbb{F}_{p^e}$ be the finite field with p^e elements. Let $0 \leq k \leq p-1$. A polynomial $f \in \mathbb{F}_{p^e}[\mathtt{x}]$ is called a *permutation polynomial* (PP) of $\mathbb{F}_{p^e}$ if the associated mapping $x \mapsto f(x)$ from $\mathbb{F}_{p^e}$ to $\mathbb{F}_{p^e}$ is a permutation of $\mathbb{F}_{p^e}$.

The premutation behaviour of the n-th reversed Dickson polynomial of the first kind $D_n(a,x)$ and its properties were explored by the author of the present paper in [1]. It was shown in [1] that to discuss the permutation property of $D_{n,k}(a,x)$, one only has to consider $a = 1$. The cases $n = p^l$, $n = p^l + 1$, and $n = p^l + 2$, where $l \geq 0$ is an integer, were discussed in [1]. The first three cases were completely explained but the last one. We list the results of the last case obtained in [1] below.

Result 1. (see [1, Remark 2.14]) Let $k = 0$ and $l = e$. Then we have

$$D_{p^e+2,0}(1,x) = \frac{1}{2}(1-4x)^{\frac{p^e+1}{2}} - x + \frac{1}{2}$$

which is a PP of $\mathbb{F}_{p^e}$ if and only if $p^e \equiv 1 \pmod 3$.

Result 2.(see [1, Theorem 2.15]) Let $k = 2$. Then $D_{p^l+2,k}(1,x)$ is a PP of $\mathbb{F}_{p^e}$ if and only if $l = 0$.

Result 3.(see [1, Theorem 2.16]) Let $p > 3$ and $k = 4$. Then $D_{p^l+2,k}(1,x)$ is a PP of $\mathbb{F}_{p^e}$ if and only if the binomial $x^{\frac{p^l-1}{2}} - \frac{1}{2}x$ is a PP of $\mathbb{F}_{p^e}$.

Result 4.(see [1, Theorem 2.17]) Let $n = p^l + 2$ and $k \neq 0, 2, 4$. Then $D_{n,k}(1,x)$ is a PP of $\mathbb{F}_{p^e}$ if and only if the trinomial $(4-k)\,x^{\frac{p^l+1}{2}} + k\,x^{\frac{p^l-1}{2}} + (2-k)x$ is a PP of $\mathbb{F}_{p^e}$.

This paper is a result of a question asked by Xiang-dong Hou. He asked the author (private communication) "when is the trinomial in result 4 a PP of $\mathbb{F}_{p^e}$?". This paper answers that question completely.

The paper is organized as follows.

In Section 2, we explain a family of permutation trinomials over $\mathbb{F}_{p^e}$ arising from the reversed Dickson polynomials when $p > 3$ is odd and k is an integer such that $k \neq 0, 2, 4$.

2010 *Mathematics Subject Classification.* 11T06, 11T55.

Key words and phrases. Finite field, Permutation polynomial, Binomial, Trinomial, Reversed Dickson polynomial.

In Section 3, we explain a family of permutation binomials over $\mathbb{F}_{p^e}$ arising from the reversed Dickson polynomials when $p = 3$.

2. A family of permutation trinomials

Theorem 2.1. *Let $p > 3$ be an odd prime and $q = p^e$, where e is a non-negative integer. Let k be an integer such that $k \neq 0, 2, 4$ and $0 \leq k \leq p-1$. Let*

$$f(x) = (4-k)x^{\frac{p^l+1}{2}} + kx^{\frac{p^l-1}{2}} + (2-k)x.$$

Then $f(x)$ is a PP of $\mathbb{F}_q$ if and only if $l = 0$ and $k \neq 3$.

Proof. Assume $l = 0$ and $k \neq 3$.

Since $l = 0$, $f(x) = 2(3-k)x + k$. Since $k \neq 3$, clearly $f(x)$ is a PP of $\mathbb{F}_q$.

Consider the following three cases.

Case 1. $l = 0$ and $k = 3$

Case 2. $l \neq 0$ and $k = 3$

Case 3. $l \neq 0$ and $k \neq 3$

Now we claim that $f(x)$ is not a PP of $\mathbb{F}_{p^e}$ in each case above.

Case 1. Since $l = 0$ and $k = 3$, $f(x) = 3$, which is clearly not a PP of $\mathbb{F}_{p^e}$.

Case 2. Let $l \neq 0$ and $k = 3$. Then

$$f(x) = x^{\frac{p^l+1}{2}} + 3x^{\frac{p^l-1}{2}} - x.$$

Note that $f(1) = 3$ and

$$f(-1) = (-1)^{\frac{p^l+1}{2}} + 3(-1)^{\frac{p^l-1}{2}} + 1.$$

which implies

$$f(-1) = \begin{cases} -1 & , \frac{p^l+1}{2} \text{ is even}, \\ 3 & , \frac{p^l+1}{2} \text{ is odd}. \end{cases}$$

Clearly, $f(x)$ is not a PP when $\frac{p^l+1}{2}$ is odd.

When $\frac{p^l+1}{2}$ is even, we have $p^l + 1 \equiv 0 \pmod 4$ and hence $p > 5$. It is clear that $f(4) \equiv 3 \pmod p$ when $\frac{p^l+1}{2}$ is even.

Hence $f(x)$ is not a PP of $\mathbb{F}_{p^e}$ in Case 2.

Case 3. Let $l \neq 0$ and $k \neq 3$. Consider

$$f(x) = (4-k)x^{\frac{p^l+1}{2}} + kx^{\frac{p^l-1}{2}} + (2-k)x.$$

Note that $f(0) = 0$. Also note that $f(\mathbb{F}_p) \subseteq \mathbb{F}_p$.

Sub Case 3.1. $l = (2n)e$, where $n \in \mathbb{Z}^+$. Then we have

$$\frac{p^l+1}{2} = \frac{p^{(2n)e}+1}{2} = \frac{(p^{ne}-1)(p^{ne}+1)}{2} + 1 \equiv 1 \pmod{p^e - 1},$$

which implies

$$f(x) = (4-k)x^{\frac{p^l+1}{2}} + kx^{\frac{p^l-1}{2}} + (2-k)x \equiv 2(3-k)x + kx^{p^e-1} \pmod{x^{p^e} - x}.$$

Since

$$2(k-3)x \equiv k \pmod{p}$$

has a non-zero solution, there exists a non-zero $x \in \mathbb{F}_p$ such that $f(0) = 0$. Hence $f(x)$ is not a PP of $\mathbb{F}_{p^e}$.

(or by Hermite's criterion, $f(x)$ is not a PP of $\mathbb{F}_{p^e}$ since the degree of $f(x) > p^e - 2$).

Sub Case 3.2. $l \neq (2n)e$, where n is a non-negative integer. Note that here we only need to consider $1 \leq l \leq 2e-1$ since

$$(4-k)x^{\frac{p^{2e+i}+1}{2}} + kx^{\frac{p^{2e+i}-1}{2}} + (2-k)x \equiv (4-k)x^{\frac{p^i+1}{2}} + kx^{\frac{p^i-1}{2}} + (2-k)x \pmod{x^{p^e} - x},$$

for $1 \leq i \leq 2e-1$.

So let $1 \leq l \leq 2e-1$ and consider

$$f(x) = (4-k)x^{\frac{p^l+1}{2}} + kx^{\frac{p^l-1}{2}} + (2-k)x.$$

Let $x \in \mathbb{F}_p^*$ and l be even. Then

$$\begin{aligned} f(x) &= (4-k)x^{\frac{p^l-1}{2}+1} + kx^{\frac{p^l-1}{2}} + (2-k)x \\ &= 2(3-k)x + k. \end{aligned}$$

Since

$$2(k-3)x \equiv k \pmod{p}$$

has a non-zero solution, there exists a non-zero $x \in \mathbb{F}_p$ such that $f(0) = 0$. Hence $f(x)$ is not a PP of $\mathbb{F}_{p^e}^*$.

Assume that l is odd. Then clearly for $x \in \mathbb{F}_{p^e} \setminus \mathbb{F}_p$, $f(x) \notin \mathbb{F}_p$.

Since l is odd, for $x \in \mathbb{F}_p$ we have

$$f(x) = (4-k)x^{\frac{p+1}{2}} + kx^{\frac{p-1}{2}} + (2-k)x$$

which implies

$$f(x)^2 = k^2 x^{p-1} + \text{ terms with lower degree.}$$

By Hermite's criterion, $f(x)$ does not permute $\mathbb{F}_p$. Hence $f(x)$ is not a PP of $\mathbb{F}_{p^e}$. □

3. A family of permutation binomials

Let $p = 3$. Since $k \neq 0, 2$, we have $k = 1$. Then

$$f(x) = (4-k)x^{\frac{p^l+1}{2}} + kx^{\frac{p^l-1}{2}} + (2-k)x = x^{\frac{p^l-1}{2}} + x.$$

Lemma 3.1. *(see [3]) Pick $d, r > 0$ with $d \mid (q-1)$, and let $h \in \mathbb{F}_q[x]$. Then $f(x) = x^r h(x^{\frac{q-1}{d}})$ permutes $\mathbb{F}_q$ if and only if*

(1) $\gcd(r, \frac{q-1}{d}) = 1$, *and*

(2) $x^r h(x)^{\frac{q-1}{d}}$ *permutes* μ_d.

Theorem 3.2. *Let $p = 3$ and $q = 3^e$, where e is a non-negative integer. Let*

$$f(x) = x^{\frac{p^l-1}{2}} + x.$$

Then $f(x)$ is a PP of $\mathbb{F}_q$ if and only if

(i) $l = 0$, *or*
(ii) $l = me + 1$, *where m is a non-negative even integer.*

Proof. When $l = 0$, $f(x) = x + 1$ which is a PP of $\mathbb{F}_q$.

Let $l = me + 1$, where m is a non-negative even integer. Note that since m is a non-negative even integer, we have

$$\frac{3^{me+1}-1}{2} = \frac{3^{me}-1}{2} + 3^{me} = \frac{(3^{\frac{me}{2}}-1)(3^{\frac{me}{2}}+1)}{2} + 3^{me} \equiv 1 \pmod{3^e-1}.$$

So, when $l = me+1$, where m is a non-negative even integer, $f(x) \equiv 2x \pmod{x^q-x}$ which is clearly a PP of $\mathbb{F}_q$.

Now assume that $l \neq 0$ and $l = me + 1$, where m is a non-negative odd integer.

$$\frac{3^{me+1}-1}{2} \pmod{3^e-1} \text{ is } \begin{cases} \text{even} & \text{, e is odd,} \\ \text{odd} & \text{, e is even.} \end{cases}$$

If $\frac{3^{me+1}-1}{2} \pmod{3^e-1}$ is even, then $f(0) = 0 = f(-1)$ which implies $f(x)$ is not a PP of $\mathbb{F}_q$.

Now consider the case where $\frac{3^{me+1}-1}{2} \pmod{3^e-1}$ is odd. Note that in this case e is even. Also,
$\frac{3^{me+1}-1}{2} \equiv \frac{3^e-1}{2} + 1 \pmod{3^e-1}$. Then

$$f(x) = x^{\frac{3^{me+1}-1}{2}} + x \equiv x^{\frac{3^e-1}{2}+1} + x \pmod{x^{3^e}-x}.$$

Now we claim that $x^{\frac{3^e-1}{2}+1} + x$ is not a PP of $\mathbb{F}_{3^e}$.

$$x^{\frac{3^e-1}{2}+1} + x = x(x^{\frac{3^e-1}{2}} + 1) = x\,h(x^{\frac{3^e-1}{2}}),$$

where $h(x) = x + 1$. $\gcd(1, \frac{3^e-1}{2}) = 1$, but $-1 \in \mu_2$ and $h(-1) = 0$. So $x\,h(x)^{\frac{3^e-1}{2}}$ does not permute μ_2. Then by Lemma 3.1, $f(x)$ is not a PP of $\mathbb{F}_{3^e}$.

Hence the proof.

□

References

[1] N. Fernando, *Reversed Dickson Polynomials of the $(k+1)$-th kind over finite fields.* arXiv:1605.04338

[2] R. Lidl and H. Niederreiter, *Finite Fields*, 2nd ed., Cambridge Univ. Press, Cambridge, 1997.

[3] M. E. Zieve, *On some permutation polynomials over $\mathbb{F}_q$ of the form $x^r h(x^{\frac{q-1}{d}})$*, Proc. Am. Math. Soc. 137 (2009), 2209 – 2216.

Department of Mathematics, Northeastern University, Boston, MA 02115
E-mail address: w.fernando@northeastern.edu